

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human DNAM-1/CD226 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human DNAM-1/CD226 Glu19-Asn247 Accession # Q15762
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human DNAM-1/CD226 Fc Chimera (Catalog # 666-DN)
Neutralization		Measured by its ability to neutralize DNAM-1-mediated adhesion of the COLO 205 human colorectal adenocarcinoma cell line. Shibuya, A. <i>et al.</i> (1996) <i>Immunity</i> 4:573. The Neutralization Dose (ND ₅₀) is typically 0.5-1.5 µg/mL in the presence of 20 µg/mL Recombinant Human DNAM-1/CD226 Fc Chimera.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

DNAX accessory molecule-1 (DNAM-1), also known as CD226, is a 65 kDa type I transmembrane glycoprotein in the immunoglobulin superfamily (1). Mature human DNAM-1 contains a 236 amino acid (aa) extracellular domain (ECD) with two Ig-like C2-set domains and a 61 aa cytoplasmic region that contains motifs for binding PDZ domains and band 4.1 family proteins (1, 2). Within the ECD, human DNAM-1 shares 50% and 52% aa sequence identity with mouse and rat DNAM-1, respectively. DNAM-1 is expressed on multiple lymphoid and myeloid cells and interacts with CD155/PVR and Nectin-2/CD112 (3, 4). Ligation of DNAM-1 promotes the activation of NK cells, CD8⁺ T cells, and mast cells (2-6), dendritic cell maturation, megakaryocyte and activated platelet adhesion to vascular endothelial cells, and monocyte extravasation; it inhibits the formation of osteoclasts (7-10). Platelet-endothelium interactions mediated by DNAM-1 enable the metastasis of tumor cells to the lung (11). In activated, but not in resting NK, T, and mast cells, the *cis* association of DNAM-1 with CD18 contributes to the tyrosine and serine phosphorylation of DNAM-1 during activation (6, 9, 12-14).

References:

1. Fuchs, A. and M. Colonna (2006) *Semin. Cancer Biol.* **16**:359.
2. Shibuya, A. *et al.* (1996) *Immunity* **4**:573.
3. Bottino, C. *et al.* (2003) *J. Exp. Med.* **198**:557.
4. Tahara-Hanaoka, S. *et al.* (2004) *Int. Immunol.* **16**:533.
5. Dardalhon, V. *et al.* (2005) *J. Immunol.* **175**:1558.
6. Bachelet, I. *et al.* (2006) *J. Biol. Chem.* **281**:27190.
7. Reymond, N. *et al.* (2004) *J. Exp. Med.* **199**:1331.
8. Kakehi, S. *et al.* (2007) *Mol. Cell. Biochem.* **301**:209.
9. Kojima, H. *et al.* (2003) *J. Biol. Chem.* **278**:36748.
10. Tahara-Hanaoka, S. *et al.* (2006) *Blood* **107**:1491.
11. Morimoto, K. *et al.* (2007) *Oncogene* July 16 epub.
12. Shibuya, K. *et al.* (1999) *Immunity* **11**:615.
13. Shibuya, K. *et al.* (2003) *J. Exp. Med.* **198**:1829.
14. Shibuya, A. *et al.* (1998) *J. Immunol.* **166**:1671.