

**DESCRIPTION**


<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects human and mouse β-1,3-N-acetylglucosaminyltransferase 1/B3GNT1 in Western blot. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) B4GalT1 and rhB3GNT 2 is observed.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human β-1,3-N-acetylglucosaminyltransferase 1/B3GNT1 Asp43-Cys415 Accession # O43505
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 μg/mL	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.2 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

B3GNT1 (β-1,3-N-acetylglucosaminyltransferase 1; also iGnT) is a 50 kD member of the β3GnT family of enzymes. It is synthesized by sensory neurons, endothelial cells and skeletal muscle. In the trans-Golgi, it collaborates with B4GALT1 to generate a linear poly-N-acetylactosamine sequence that can be incorporated into either N-linked or O-linked glycans. Human B3GNT1 is a 451 amino acid (aa) type II transmembrane protein. It contains an eight aa N-terminal cytoplasmic segment, an extended 28 aa transmembrane region (aa 9-36), and a 379 aa luminal domain (aa 37-415). Over aa 37-415, human B3GNT1 shares 97% aa identity with mouse B3GNT1.