

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Fcε RIα in direct ELISAs.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Fcε RIα Val26-Gln205 Accession # NP_001992
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The α subunit of the high affinity IgE receptor (Fcε RIα or Fcε R1A) is an IgE-binding type I transmembrane glycoprotein of the multichain immune recognition (MIRR) family (1, 2). The receptor, Fcε RI, is a tetrameric complex of one α, one β and two γ subunits (αβγ₂) on mast cells and basophils (1). An alternate trimeric form (αγ₂) is expressed on human, but not rodent, mast cells, basophils, eosinophils and professional antigen presenting cells (3). While the γ subunit is essential for expression of Fcε RIα on the cell surface and for cell signaling, the β subunit, when present, increases the half-life of the Fcε RI complex on the cell surface (3, 4). An isoform of the β subunit, βT, blocks processing of the α subunit and its cell surface expression (2, 3, 5). Human Fcε RIα cDNA encodes 257 amino acids (aa) including a 25 aa signal sequence, a 180 aa extracellular domain containing two Ig-like domains that bind IgE and an endoplasmic reticulum retention motif, a 21 aa transmembrane domain with a charged amino acid (Asp219) that contributes to intracellular transport, and a 32 aa cytoplasmic sequence (1, 3, 6). Human Fcε RIα shares 50-62% aa sequence identity with mouse, rat, equine, ovine, bovine, porcine and canine Fcε RIα. Binding of IgE alone increases surface expression of Fcε RI, while crosslinking of IgE/Fcε RI complexes by IgE ligands (allergens) initiates receptor internalization and signaling (2, 4, 5). Mast cell and basophil activation by IgE/Fcε RI crosslinking causes degranulation, releasing histamine, leukotrienes, prostaglandins, and other mediators of immediate-type and late-phase allergic reactions. Circulating autoantibodies that crosslink Fcε RIα are often found in patients with chronic urticaria (7). Fcε RI on human antigen presenting cells mediates uptake and processing of allergens for presentation by class II MHC (2, 3). Fcε RI expression on human DC and Langerhans cells is up-regulated during allergic reactions (atopy) and correlates with serum IgE concentration (3).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.