

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human α -N-acetylgalactosaminidase/NAGA in direct ELISAs and Western blots. In direct ELISAs, less than 2% cross-reactivity with recombinant human α -Galactosidase A is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line recombinant human α -N-acetylgalactosaminidase/NAGA Leu18-Gln411 Accession # P17050
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

NAGA is a lysosomal α -N-acetylgalactosaminidase that cleaves non-reducing α -N-acetylgalactosaminyl moieties from glycoconjugates (1). Mature NAGA has 394 amino acids and is trafficked to the lysosome via the mannose-6-phosphate receptor-mediated pathway (2). The enzyme is a retaining exoglycosidase, where both the substrate and product of the enzymatic reaction have the same anomeric configuration (3). Deficiency in NAGA results in increased urinary excretion and tissue accumulation of glycopeptides and oligosaccharides containing terminal α -N-acetylgalactosaminyl moieties (4), manifesting as Schindler's disease, an autosomal recessive disease with neuroaxonal dystrophy and other neurological symptoms (5). The enzyme can be used to remove α -N-acetylgalactosaminyl residues present on red blood cells thus converting blood type A to blood type O (6, 7, 8).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.