

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Thymosin β 4 in direct ELISAs and Western blots. In direct ELISAs, approximately 5% cross-reactivity with recombinant human (rh) Thymosin β 10 and rhThymosin β 16 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Thymosin β 4 Ser2-Ser44 Accession # P62328
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Thymosin beta 4 (T β 4; also TB4X and Fx) is a 5.0 kDa member of the β -thymosin family of molecules. Members of this family range from 41-44 amino acids (aa) in length, and possess an isoelectric point that lies between pH 4.0-7.0 (α -thymosins have values less than 4.0). Multiple cell types produce T β 4, either constitutively, or after stimulation. They include platelets, endothelial cells, neutrophils, astrocytes and macrophages. T β 4 is both a secreted and intracellular molecule. The secreted form contributes to wound healing and angiogenesis, and may act on ATPase. Intracellularly, it forms a 1:1 complex with G-actin and blocks F-actin polymerization. This regulates the availability of actin monomers for filament formation and subsequent cell migration. Mature human T β 4 is 43 aa in length (aa 2-44). It contains an actin-binding site (aa 17-23), one acetylated Ser and five acetylated lysines (4; 12; 26; 32; 39) and one phosphorylation site at Thr23. T β 4 undergoes proteolytic processing to generate an N-terminal acetylated peptide (aa 2-5: SerAspLysPro). Mature human T β 4 is identical to mouse T β 14 in aa sequence, and it shares 74% aa identity with its human family member T β 10.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.