

DESCRIPTION

| | |
|---|---|
| Species Reactivity | Mouse |
| Specificity | Detects mouse CD79B in direct ELISAs and Western blots. In direct ELISAs, approximately 55% of cross-reactivity with human CD79B is observed. |
| Source | Polyclonal Sheep IgG |
| Purification | Antigen Affinity-purified |
| Immunogen | Chinese hamster ovary cell line CHO-derived recombinant mouse CD79B Met29-Asp158 Accession # P15530 |
| Conjugate | Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm |
| Formulation | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide |
| *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. | |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

| | |
|----------------------------|--|
| CyTOF-ready | Optimal dilution of this antibody should be experimentally determined. |
| Western Blot | Optimal dilution of this antibody should be experimentally determined. |
| Flow Cytometry | Optimal dilution of this antibody should be experimentally determined. |
| Immunocytochemistry | Optimal dilution of this antibody should be experimentally determined. |

PREPARATION AND STORAGE

| | |
|--------------------------------|---|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied |

BACKGROUND

CD79B (also known as B29, Igβ and B cell antigen receptor complex-associated protein β-chain) is a 37-39 kDa member of the Ig-Superfamily. It is expressed on B cells, and forms a covalent heterodimer with CD79a. This complex interacts noncovalently with membrane Ig, forming the B cell antigen receptor. Within this complex, membrane Ig detects antigen while CD79a:b initiates signaling. CD79B is also required for formation of pre-B cells during B cell development. Mature mouse CD79B is a 203 amino acid (aa) type I transmembrane glycoprotein (aa 26-228). It contains an extracellular region with one V-type Ig-like domain (aa 41-132) and an ITAM-containing cytoplasmic domain (aa 181-228). CD79B may migrate as two bands in SDS-PAGE. One defines the standard 37 kDa form, while a second represents one of two possible isoforms, the first of which is an underglycosylated full-length CD79B, and the second of which is an alternative splice form that likely lacks the C-terminal 32 amino acids. Mouse CD79a and CD79B share only 24% aa identity. Over aa 29-158, mouse CD79B shares 54% and 78% aa identity with human and rat CD79B, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.