

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-4 Rα in direct ELISAs. In direct ELISAs, approximately 10% cross-reactivity with recombinant mouse IL-4 R is observed, and less than 1% cross-reactivity with recombinant human (rh) IL-5 Rα, rhIL-5 Rβ, rhIL-9 R, rhIL-13 Rα1, and rhIL-13 Rα
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human IL-4 Rα Gly24-His232 Accession # P24394
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 4 (IL-4) is a pleiotropic cytokine produced by activated T cells, mast cells, and basophils. The biological functions of IL-4 are mediated by the binding of IL-4 to high-affinity cell surface receptor complexes. Two types of IL-4 receptor complexes have been described. The type I IL-4 receptor complex is composed of a high-affinity IL-4-binding subunit (referred to as IL-4 Rα) and the common γ chain that does not bind IL-4 by itself. The type II IL-4 receptor complex is composed of IL-4 Rα and IL-13 Rα1. Besides IL-4 signals, the type II IL-4 receptor complex can also transduce IL-13 signals. In the type II complex, the IL-4 Rα subunit binds only IL-4 and not IL-13. Similarly, the IL-13 Rα1 subunit binds only IL-13 and not IL-4. The cDNA clones for both the human and mouse IL-4 Rα have been isolated and shown to encode an approximately 140 kDa type I transmembrane protein with a large cytoplasmic domain that is essential for signal transduction. In mouse cells, an alternatively spliced variant encoding a soluble secreted IL-4 Rα isoform has also been identified. Naturally occurring soluble IL-4 Rα that binds IL-4 with high-affinity has been found in mouse and human biological fluids.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.