

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human MTF2 in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human MTF2 Asn356-Ser554 Accession # Q9Y483
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

MTF2 (Metal-response element-binding transcription factor 2; also M96, ZIRF1 and PCL2) is a 67 kDa member of the polycomb group of proteins. It appears to participate in the activities of at least two complexes, PRC1 (Polycomb repressive complex 1) and PRC2. In general, the PRC complexes are repressive. In stem cells, MTF2/PRC2 inhibits expression of pluripotency factors such as KLF4 and Tbx3, likely by recruiting the full PRC2 complex to target gene promoters. And during AP specification of the axis, MTF2, as part of both a PRC1 and PRC2 complex, blocks Hox gene expression. Human MTF2 is 593 amino acids (aa) in length. It contains one Tudor domain (aa 40-91), two PHD-type Zn-finger regions (aa 102-157 and 201-255) and three utilized phosphorylation sites at Ser24, 27 and 488. There are multiple potential splice variants. One is 60 kDa in size and shows a deletion of aa 331-387, while a second is 55 kDa in size and possesses an alternative start site at Met103. A third contains a combination of the above, a fourth possesses a 34 aa substitution for aa 1-307, and a fifth utilizes an alternative start site at Met332. Over aa 356-554, human MTF2 shares 94% aa identity with mouse MTF2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.