

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FoxP4 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) FoxP1, rhFoxP2, rhFoxP3, and rhCTBP1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human FoxP4 Pro549-Ser680 Accession # Q8IVH2
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

FoxP4 (Forkhead box protein P4; also FKHLA) is a 70-80 kDa member of the winged helix transcription factor gene family, FoxP subfamily of proteins. It is a transcriptional repressor that is strongly expressed in embryonic gut and lung epithelium, and in neurons both pre-and postnatally. Postnatally, it is found in cerebellar Purkinje cells. During development, it is found in hippocampal hilus, proliferating cells of the ventricular zone, and neurons of the basolateral nucleus of the amygdala. Human FoxP4 is 680 amino acids (aa) in length. It contains a large poly-Gln region (aa 65-219), a C2H2 Zn-finger domain (aa 307-332), a Leu-zipper dimerization motif (aa 349-370) and a forkhead DNA binding domain (aa 467-559). Phosphorylation occurs at Ser360. FoxP4 forms homodimers, and heterodimers with FoxP1 and FoxP2; it does not bind CTBP1. One potential splice form shows a deletion of aa 384-395. Over aa 549-680, human and mouse FoxP4 share 89% aa identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.