

DESCRIPTION

Species Reactivity	Mouse/Rat
Specificity	Detects mouse and rat Plexin B3 in direct ELISAs and Western blots. In direct ELISAs, approximately 38% cross-reactivity with recombinant human (rh) Plexin B3 is observed, and less than 1% cross-reactivity with rhPlexin B2 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse Plexin B3 His25-Gln1235 Accession # Q9QY40
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Plexin-B3 (also plexin-6) is a 260-270 kDa member of the plexin-B subfamily, plexin family of proteins. It is found on postnatal neurons and oligodendroglia, and appears to have opposing functions. In a context-dependent manner, plexin-B3 may block cell process extension and induce growth cone collapse by binding to Sema5. Conversely, it may engage in homophilic interactions *in trans*, promoting growth cone extension and axon elongation. The HGF receptor Met has been reported to complex with plexin-B3 *in cis*, and this interaction may contribute to its reported opposing effects. Mature mouse plexin-B3 is an 1868 amino acid (aa) type I transmembrane glycoprotein. It contains a 1211 aa extracellular domain (ECD) (aa 25-1892) plus a 636 aa cytoplasmic region. The ECD contains one Sema-domain (aa 25-451), two PSI domains (aa 454-646) and four IPT regions (aa 814-1224). There is one isoform variant that contains an alternative Met start site 10 aa upstream of the standard site. Based on human studies, proteolysis of mouse plexin-B3 likely occurs, generating 160 kDa and 140 kDa ECD fragments. Over aa 25-1235, mouse plexin-B3 shares 95% and 82% aa identity with rat and human plexin-B3, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.