

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human RGS8 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human RGS4 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human RGS8 Asn10-Thr76 Accession # P57771
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

RGS8 (Regulator of G-protein signaling 8) is a 22-23 kDa (predicted) member of the B/R4 subfamily, RGS family of proteins. It is expressed in NK cells and neurons, particularly cerebellar Purkinje cells, and is known to regulate GPCR signaling. When GPCRs are activated, a GPCR cytoplasmic domain associated Gαβγ heterotrimer dissociates in response to GTP binding. This generates Gα:GTP and Gβγ dimers that initiate downstream signaling. Gα has GTPase activity, and when GTP is hydrolyzed, the Gαβγ trimer reforms and stops signaling. RGS8 potentiates Gα GTP hydrolase activity, particularly when associated with M1 muscarinic acetylcholine receptors. Human RGS8 is 180 amino acids (aa) in length, and is characterized by the presence of one RGS domain (aa 56-171), and an N-terminal GPCR binding motif (aa 6-9). There is one splice variant that shows a 27 aa substitution for aa 1-9. Expression of this variant may negatively impact its regulation of Gα-coupled responses. Over aa 10-76, human RGS8 shares 98% aa identity with mouse RGS8.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.