

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human XCL1/Lymphotactin in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human XCL1/Lymphotactin Val22-Gly114 Accession # P47992
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human lymphotactin (Lptn)/XCL1 (also named human SCM-1α and ATAC) and its mouse homologue belong to the C or γ subfamily of chemokines. The C chemokines lack two (the 1st and 3rd) of the four invariant cysteine residues normally found in the CC and CXC chemokines and have an extended carboxy terminus. Human lymphotactin encodes a 114 amino acid residue precursor protein with a 21 amino acid residue predicted signal peptide. The expression of lymphotactin is abundant in some activated T cells such as activated CD8⁺ T cells and other class I MHC restricted T cells. Lptn expression is absent in CD4⁺ T cells. Human and mouse Lptn share approximately 60% amino acid sequence homology. The gene for lymphotactin has been mapped to chromosome 1 in both human and mouse. Recombinant human lymphotactin has been shown to have chemotactic activity for lymphocytes and NK cells. The orphan receptor GPR5 has been reported to be the specific receptor for Lptn.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.