

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects mouse EBF-2 in direct ELISAs. Detects mouse and human EBF-2 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse EBF-2 Arg407-Ser519 Accession # O08792
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

EBF-2 (Early B cell Factor 2; also Mmot1, OLF3 and COE2) is a 62 kDa (predicted) member of the COE family of transcription factors. It is expressed in immature osteoblasts and Purkinje cells, and in the embryo is associated with the migration of postmitotic neuroblasts. In immature osteoblasts, EBF-2 appears to upregulate OPG, suppressing osteoclast formation. And in the developing retina, EBF-2 is found in ganglion, glycinergic Amacrine and horizontal cells, possibly promoting their development over that of photoreceptor cells. Mouse EBF-2 is 575 amino acids (aa) in length. It contains one DNA-binding region with an embedded C5-type zinc-finger motif (aa 62-238), a dimerization ITP/TIG domain (aa 253-336), and a Pro/Ser-rich transactivation domain (aa 453-534). Although considered an HLH type transcription factor, it does not contain the typical "b", or basic amino acid sequence associated with bHLH factors. EBF-2 both homodimerizes, and heterodimerizes with EBF-1 and -3. There is an alternative start site at Met23. Over aa 407-519, mouse EBF-2 is identical in aa sequence to rat EBF-2 and shares 99% aa identity with human EBF-2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.