

DESCRIPTION

Species Reactivity	Mouse/Rat
Specificity	Detects recombinant mouse Inositol Monophosphatase 3/IMPAD1 in direct ELISAs and Western blots. Detects mouse Inositol Monophosphatase 3/IMPAD1 and rat Inositol Monophosphatase 3/IMPAD1 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse Inositol Monophosphatase 3/IMPAD1 Asn145-His356 Accession # Q80V26
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IMPAD1 (Inositol monophosphatase domain-containing protein 1; also IMPA3, gPAPP and IMPase 3) is a 40-42 kDa member of the inositol monophosphatase family of proteins. It is expressed in embryo, and found in Purkinje cells, brain stem, lung and chondrocytes. IMPAD1 in theory may catalyze the synthesis of myo-inositol from myo-inositol monophosphate. Free myo-inositol is used to generate inositol phospholipid, an essential component of intracellular signaling pathways that mobilize calcium. IMPAD1 is reported to promote sulfation of chondroitin by converting PAP (an endproduct of the sulfation process) to 5'-AMP. PAP is a known inhibitor of SULTs. IMPAD1 is a 356 amino acid (aa) type II transmembrane Golgi-embedded glycoprotein. It contains a short cytoplasmic tail (aa 1-12) and an extended luminal region (aa 34-356) that contains its catalytic domain (aa 60-347). Over aa 145-356, mouse IMPAD1 shares 99% and 93% aa sequence identity with rat and human IMPAD1, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.