

**DESCRIPTION**

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human GM-CSF R $\alpha$ in direct ELISAs and Western blots.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human GM-CSF R $\alpha$ Extracellular domain
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.1 $\mu$ g/mL	Recombinant Human GM-CSF R $\alpha$ (Catalog # 706-GR)
<b>Flow Cytometry</b>	0.25 $\mu$ g/10 <sup>6</sup> cells	Human whole blood monocytes
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Granulocyte macrophage colony stimulating factor receptor alpha (GM-CSF R $\alpha$ ), also known as CD116, is a component of the receptor complex that mediates cellular responses to GM-CSF. GM-CSF promotes the differentiation and mobilization of granulocyte-macrophage, erythroid, megakaryocyte, and eosinophil progenitors. It enhances the activation of myeloid cell effector functions and plays a role in the development of Th1 biased immune responses, allergic inflammation, and autoimmunity (1-4). Mature human GM-CSF R $\alpha$  is an 80 kDa type I transmembrane glycoprotein that consists of a 298 amino acid (aa) extracellular domain (ECD) with two fibronectin type III domains and a juxtamembrane WSXWS motif, a 26 aa transmembrane segment, and a 54 aa cytoplasmic domain (5). Within the ECD, human GM-CSF R $\alpha$  shares approximately 33% aa sequence identity with mouse and rat GM-CSF R $\alpha$ . Alternate splicing of human GM-CSF R $\alpha$  generates several additional isoforms that lack the cytoplasmic and/or transmembrane regions. Soluble forms of the receptor retain the ability to bind GM-CSF (6, 7). GM-CSF R $\alpha$  is expressed on hematopoietic stem cells, progenitor and differentiated cells in the myeloid lineage, vascular endothelial cells, placenta, and non-hematopoietic solid tumor cells (8). GM-CSF R $\alpha$  associates with the common beta chain/CD131 ( $\beta_c$ ), a 135 kDa transmembrane protein that is also the signal transducing component of the receptors for IL-3 and IL-5 (9, 10). Association with  $\beta_c$  converts GM-CSF R $\alpha$  from a low affinity to a high affinity receptor for GM-CSF (9-11). The shared usage of  $\beta_c$  underlies the synergism between GM-CSF, IL-3, and IL-5 in their effects on myeloid cell differentiation and activation (1, 2).

**References:**

1. Martinez-Moczygamba, M. and D.P. Huston (2003) J. Allergy Clin. Immunol. **112**:653.
2. Fleetwood, A.J. *et al.* (2005) Crit. Rev. Immunol. **25**:405.
3. Eksioglu, E.A. *et al.* (2007) Exp. Hematol. **35**:1163.
4. Cao, Y. (2007) J. Clin. Invest. **117**:2362.
5. Gearing, D.P. *et al.* (1989) EMBO J. **8**:3667.
6. Pelley, J.L. *et al.* (2007) Exp. Hematol. **35**:1483.
7. Raines, M.A. *et al.* (1991) Proc. Natl. Acad. Sci. **88**:8203.
8. Chiba, S. *et al.* (1990) Cell Regul. **1**:327.
9. Kitamura, T. *et al.* (1991) Proc. Natl. Acad. Sci. **88**:5082.
10. Hayashida, K. *et al.* (1990) Proc. Natl. Acad. Sci. **87**:9655.
11. Hoang, T. *et al.* (1993) J. Biol. Chem. **268**:11881.