

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CEBP α in direct ELISAs. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) CEBP β , rhCEBP γ , rhCEBP δ , rhCEBP ζ , and rhCEBP ϵ is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CEBP α Met1-Ala124 Accession # P49715
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CEBP alpha (CCAAT/Enhancer-Binding Protein; also C/EBP) is a 42-43 kDa member of the C/EBP subfamily, bZIP family of proteins. It is expressed in multiple cell types, particularly those that are fully differentiated, and includes hepatocytes, skeletal muscle cells, adipocytes, intestinal epithelium and proliferating macrophages. CEBP α acts as a transcriptional activator for a large and diverse number of genes, including glycogen synthase, GLUT4, and G-CSFR, and participates in both cell cycle progression and differentiation. Human CEBP α is 358 amino acids (aa) in length. It contains one transactivation domain (aa 1-96), poly-Gly segment (aa 99-104), a poly-Pro motif (aa 183-189), a bZIP dimerization region (aa 281-334), and an overlapping DNA-binding Leu-zipper domain (aa 317-345). CEBP α forms both homodimers, and heterodimers with other subfamily members. One alternative start site at Met120 generates a low activity 30-32 kDa isoform variant. There is also a utilized phosphorylation site at Ser21 that regulates activity. Over aa 1-124, human CEBP α shares 93% aa identity with mouse CEBP α .

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.