

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Hormone-sensitive Lipase/HSL in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Hormone-sensitive Lipase/HSL Met302-Leu425 Accession # Q05469
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

LIPE (also known as hormone sensitive lipase/HSL) is a 88-90 kDa member of the GDXG lipolytic enzyme family of molecules. It is expressed in multiple cell types, including skeletal muscle, adrenal gland and adipocytes, and is regulated by a series of phosphorylations. Catecholamines activate the enzyme via PKA through the phosphorylation of Ser853, Ser855, and Ser951, while insulin depresses its activity via PDE3B. LIPE acts on triglycerides to release free fatty acids, and serves as a retinyl ester hydrolase. Human LIPE is 1076 amino acids (aa) in length, and contains one HSL domain (aa 302-616). This restricted 116 kDa isoform is found in testis (spermatids). There are additional isoform variants. One is 88-90 kDa in size in SDS-Page and possesses an alternative start site to the long form at Met302. This is considered the standard and most common LIPE isoform. A second is approximately 93 kDa in size and contains a 39 aa substitution for aa 1-294. A third isoform is 80 kDa in size and shows a 76 aa deletion in the center of the molecule, reducing its enzymatic activity. It is suggested that LIPE may act as both a monomer and homodimer. Over aa 302-425, human HSL shares 92-93% aa identity with both mouse and rat LIPE.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.