

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Contactin-3 in direct ELISAs. In direct ELISAs, approximately 10% cross-reactivity with recombinant mouse Contactin-3 is observed, and less than 5% cross-reactivity with recombinant human (rh) Contactin-4 and rhContactin-5 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Contactin-3 Pro26-Pro992 Accession # Q9P232
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Contactin-3, also known as BIG-1, CNTN3 and PANG, is a 150 kDa member of the TAG/F3 subgroup of Contactin neuronal adhesion proteins (1, 2). Mature human Contactin-3 consists of six immunoglobulin-like domains, a flexible linker region, four fibronectin type III domains, and a GPI anchor (3, 4). Human Contactin-3 shares 92% aa sequence identity with mouse and rat Contactin-3. Contactin-3 is primarily expressed in the brain in discrete regions including the frontal lobe, occipital lobe, cerebellum, amygdala, and hippocampus (5, 6). Alternative splicing in rat may generate a soluble isoform (4). Mouse plasmacytoma cell lines produce multiple transcripts of Contactin-3 due to the presence of IAP proviruses (3). Immobilized Contactin-3 promotes neurite outgrowth from rat hippocampal neurons (4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.