

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse HSF1 in direct ELISAs. In direct ELISAs, approximately 6% cross-reactivity with recombinant mouse HSF4 is observed and less than 1% cross-reactivity with recombinant human HSF2 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse HSF1 Thr124-Tyr247 Accession # P38532
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

HSF1 (Heat Shock Factor 1; also HSTF 1) is a 67-75 kDa member of the HSF family of proteins. It is constitutively expressed and functions as a transcriptional activator under heat shock conditions. Normally, it exists as an inactive, monomeric member of the HSP multichaperone complex. Following stress, it dissociates from the complex, homotrimerizes, undergoes multiple phosphorylations, and translocates to the nucleus. Nuclear HSF1 is SUMOylated on Lys298 and binds to heat shock elements in select genes such as HSP70. Mouse HSF1 is 525 amino acids (aa) in length. It contains a DNA binding domain (aa 15-120), two hydrophobic oligomerization regions (aa 130-203 and 380-405), a regulatory domain (aa 221-310), and a transactivation domain (aa 367-525). There are at least 13 potential Ser/Thr phosphorylation sites. When highly phosphorylated, HSF1 appears to run at about 82-85 kDa in SDS-PAGE; when SUMOylated, it runs anomalously at around 170 kDa in SDS-PAGE. One splice variant exists that shows a deletion of aa 413-434. Over aa 124-247, mouse HSF1 shares 94% and 99% aa identity with human and rat HSF1, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.