

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Amnionless in direct ELISAs and Western blots. In direct ELISAs, approximately 10% cross-reactivity with recombinant human Amnionless is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse Amnionless Ala20-Gly362 Accession # Q99JB7
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

AMN (Amnionless) is a 45-49 kDa glycoprotein that is expressed by three cell types in mouse; extraembryonic visceral endoderm, proximal tubule renal epithelium, and absorption intestinal epithelium. AMN is required for gastrulation in rodent (but not human), and participates in Vitamin B12 absorption in the gut. In particular, AMN forms a membrane complex with Cubilin, promoting its proper orientation in the membrane and subsequent binding to the Intrinsic factor: VitB12 complex. Mature mouse AMN is a 439 amino acid (aa) type I transmembrane protein. It contains a 343 aa extracellular region (aa 20-362) that possesses one VWFC domain (aa 203-254), and a 75 aa cytoplasmic tail. In human, AMN may undergo cleavage to generate a 35 kDa soluble fragment. Over aa 20-362, mouse AMN shares 92% and 67% aa sequence identity with rat and human AMN, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.