

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TLE2 in direct ELISAs. In direct ELISAs, less than 2% cross-reactivity with recombinant human (rh) TLE1, rhTLE3, and rhTLE4 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human TLE2 Ile11-Ser193 Accession # Q04725
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	5-15 µg/mL	See Below

DATA

Immunohistochemistry



TLE2 in Human Lung Mesothelioma.
TLE2 was detected in immersion fixed paraffin-embedded sections of human lung mesothelioma using Sheep Anti-Human TLE2 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF7216) at 10 µg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Sheep HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS019) and counterstained with hematoxylin (blue). Specific staining was localized to nuclei. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

TLE2 (Transducin-Like Enhancer of Split 2; also ESG2 and Grg-2) is an 83-84 kDa member of the WD repeat Groucho/TLE family of transcriptional repressors. It is ubiquitously expressed, and is known to antagonize TCF (Wnt)-mediated signaling. TLE2 binds to other family members as a heterooligomer, or to itself as a homooligomer. While it possesses no intrinsic DNA-binding activity, it does modulate the activity of multiple factors such as FoxG1, Arx and histone H3. Human TLE2 is 743 amino acids (aa) in length. It contains a Gln-rich region that mediates oligomerization (aa 1-133), a CCN domain that contains an NLS (aa 195-256), and six WD repeats that mediate protein-protein interaction (aa 455-742). There are at least five potential phosphorylation sites that if used, may account for SDS-Page MWs exceeding 95 kDa. Potential isoform variants exist. One possesses a 21 aa substitution for aa 1-9, while a second shows the same substitution coupled to an additional 10 aa substitution for aa 683-743. A third isoform possesses an alternative start site at Met56 coupled to a deletion of aa 124-190. Over aa 11-193, human TLE2 shares 91% aa identity with mouse TLE2.