

**DESCRIPTION**


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse TIGIT in direct ELISAs. In direct ELISAs, less than 2% cross-reactivity with recombinant human TIGIT is observed.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse TIGIT Gly26-Thr143 Accession # NP_001139797
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.2 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

TIGIT (T cell Immunoreceptor with Ig and ITIM domains; also Vstm3 and Vsig9) is a 30-34 kDa (in human) member of the CD28 family, Ig superfamily of molecules. It is expressed by NK cells and multiple subsets of mature T cells, and binds to PVR/CD155 and PVR2/CD112 that appear on dendritic cells (DC) and endothelium. Along with CD226, the TIGIT:CD226/DNAM1 and PVR:PVR2 pairings appear to form a network that parallels the well-characterized B7-1:B7-2 and CD28:CTLA4 system. Binding of TIGIT by DC induces DC IL-10 release and inhibits IL-12 production. Ligation of TIGIT on T cells dampens TCR-mediated activation, while NK cell TIGIT ligation blocks NK cell cytotoxicity. Mature mouse TIGIT is a type I transmembrane protein 215 amino acids (aa) in length. It contains a 116 aa extracellular region (aa 26-141) with a V-type Ig-like domain (aa 27-125), and a 79 aa cytoplasmic domain with one ITIM motif. There is one isoform variant that is quite unusual and shows an addition of nine amino acids spread over three insertion sites (SwissProt #:P86176). Mouse and human TIGIT are highly divergent, and over aa 26-143, mouse TIGIT shares only 68% aa sequence identity with human TIGIT.