

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CELSR3 in direct ELISAs. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) CELSR1 and rhCELSR2 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CELSR3 Tyr531-Ser711 Accession # Q9NYQ7
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CELSR3 (Cadherin EGF LAG Seven-pass G-type Receptor 3; also cadherin family member 11/CDHF11, FMI1 and EGFL1) is a 355 kDa (predicted) member of the LN-7TM subfamily, GPCR 2 family of proteins. It is coexpressed with Fzd3 on postmigratory neurons of the developing DRG and cranial ganglia, and appears to serve as an axonal guidance cue. Mature human CELSR3 is 3280 amino acids (aa) in length (aa 33-3312). It is a highly complex 7-transmembrane protein that contains a 2508 aa extended N-terminal extracellular region (aa 33-2540) plus a 538 aa C-terminal cytoplasmic domain. The N-terminal region contains nine consecutive cadherin domains (aa 326-1265), followed by a mixture of eight EGF-like and three laminin-like domains, and one membrane-proximal GPS domain. There are two potential isoform variants. One shows a five aa insertion after Gly2158, while another possesses an alternative start site at Met276. Over aa 531-711, human CELSR3 shares 97% aa sequence identity with mouse CELSR3.