

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse FCRL1/FcRH1 in direct ELISAs. In direct ELISAs, less than 3% cross-reactivity with recombinant human FCRL1/FcRH1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse FCRL1/FcRH1 Ala17-Ser219 Accession # Q8R4Y0
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

FCRL1 (Fc receptor-like protein 1; also CD307a, FcRH1 and IFGP1) is a 36 kDa (predicted) member of the Ig Superfamily. In mouse, it is found on mature follicular and marginal B cells, splenic T cells and NK cells. FCRL1 is likely not a receptor for immunoglobulin. At present, it would appear that mouse FCRL1 serves as an inhibiting coreceptor, in as much as it contains a series of cytoplasmic ITIM motifs. Mature mouse FCRL1 is a 327 amino acid (aa) type I transmembrane protein (aa 17-343). It contains a 203 aa extracellular region (aa 17-219) that shows two C2-type Ig-like domains (aa 17-200), and a 103 aa cytoplasmic domain. There are multiple potential splice variants. One shows a deletion of aa 206-248 (potentially soluble), a second contains a six aa substitution for aa 319-343, a third possesses a three aa substitution for aa 209-343, and a fourth shows a 21 aa substitution for Ala11. Over aa 17-219, mouse FCRL1 shares 83% aa sequence identity with rat FCRL1. Over the same region, it shares 63% aa sequence identity with human FCRL1 when a third Ig-like domain that is present in human is excluded from the analysis.