

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse, rat STI1 in Western blots and detects recombinant human STI1 in direct ELISAs.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human STI1 Met1-Arg543 Accession # P31948
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

STI1 (Stress-induced phosphoprotein 1; also HOP and STIP-1) is a 60-62 kDa member of the STI-1 family of proteins. It is widely expressed, and participates in the formation of functional steroid hormone receptor complexes. Steroid receptors (SR) require proper folding to achieve optimal conformation. In the case of progesterone and glucocorticoid receptors, interaction with chaperones hsp70 and hsp90 facilitates this folding. STI1 is not a chaperone, but a mediator of SR-hsp70-hsp90 interaction. STI1 first binds dimeric hsp90. This trimer then interacts with a SR-(monomeric)hsp70-ADP protein heterodimer to form a SR-hsp70-STI1-hsp90 pentamer that subsequently dissociates into a mature hsp90-SR complex. The STI1-hsp90 complex is also found extracellularly and may contribute to MMP-2 activation. Human STI1 is 543 amino acids (aa) in length. It contains an initial N-terminal TPR domain (aa 4-105) that binds to hsp70, followed by a second TPR domain (aa 225-461) that binds to dimeric hsp90. There are at least 10 acetylated Lys residues plus four utilized phosphorylation sites. Zone potential isoform variant is reported that contains a 48 aa substitution for Gln3. Full-length human STI1 shares 97% aa sequence identity with mouse STI1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.