

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Laminin α 4 in direct ELISAs and Western blots. In direct ELISAs, approximately 15% cross-reactivity with recombinant mouse Laminin α 4 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Laminin α 4 Gln826-Ala1816, predicted Accession # EAW48267
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Laminin α 4 (LAMA-4 [Laminin subunit alpha 4]; also Laminin-14/423, -8/411 and -9/421 subunit alpha) is a 180-230 kDa secreted member of the laminin family of molecules. It is found in the basement membranes of adipocytes, endothelial cells, cardiac and visceral smooth muscle cells, fibroblasts, keratinocytes, and pancreatic acinar cells. Laminin is a heterotrimeric glycoprotein that attaches cells to basement membranes. It is composed of covalently-linked α -, β - and γ -chains, and is known to attach cells to basement membranes. In the case of the α -subunit, this is mediated through binding to multiple integrins (α 3 β 1, α v β 3 and α 6 β 1), plus dystroglycan and the syndecans-2 and -4. Mature human Laminin α 4 is 1799 amino acids (aa) in length (aa 25-1823). It contains multiple domains, including four EGF-like domains (aa 82-255) and five Laminin G-like domains (aa 833-1820). There is a chondroitin sulfate attachment at the N-terminus that accounts for 20-30 kDa of MW. Laminin α 4 is cleaved between the Laminin G-like 3 and 4 domains, creating a soluble 44 kDa fragment that possesses antibacterial activity. There are two potential splice variants, one that shows a deletion of aa 266-272, and another that contains a 55 aa substitution for aa 66-1823. Over aa 826-1816, human Laminin α 4 shares 91% aa sequence identity with mouse Laminin α 4.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.