

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse CBFB in Western blots and recombinant human CBFB in direct ELISAs. In direct ELISAs, approximately 5% cross-reactivity with recombinant human RUNX-1, -2, and -3 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CBFB Pro2-Glu165 Accession # Q13951
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CBFB (Core binding factor beta; also PEA2β and PEBP2β) is a widely-expressed 21-24 kDa member of the CBFB family of proteins. It forms dimeric transcriptional complexes with multiple molecules, including CBFB, RUNX1 and RUNX2. Although it does not bind DNA, it potentiates the binding of its partners to DNA. Human CBFB is 182 amino acids (aa) in length. It contains multiple α-helices and β-strands. CBFB has at least two splice variants. One is approximately 23 kDa in size and possesses a 22 aa substitution for aa 166-182. A second is 16 kDa in size and contains a 22 aa substitution for aa 134-182. And a third shows a potential deletion of aa 56-94 coupled to the above 22 substitution for aa 166-182. CBFB is known to form a 68-70 kDa fusion protein with smooth muscle myosin heavy chain in AML. The CBFB contribution to the N-terminus of this fusion protein usually involves aa 1-165. Over aa 1-165, human CBFB shares 98% aa sequence identity with mouse CBFB.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.