

DESCRIPTION

Species Reactivity	Canine
Specificity	Detects canine IL-10 in direct ELISAs and Western blots. In direct ELISAs, approximately 30% cross-reactivity with recombinant feline IL-10, recombinant human IL-10, recombinant porcine IL-10 is observed and 5% cross-reactivity with recombinant mouse
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant canine IL-10 (R&D Systems, Catalog # 735-CL) Ser20-Ile179 Accession # XP_855560
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-10, initially designated cytokine synthesis inhibitory factor (CSIF), was originally identified as a product of murine T helper 2 (Th2) clones that inhibited the cytokine production by Th1 clones. The canine homologue of mouse IL-10 has been cloned. Canine IL-10 shares 80% amino acid sequence homology with human IL-10 and 72% sequence homology with mouse IL-10.

IL-10 is a pleiotropic cytokine that can exert either immunostimulatory or immunosuppressive effects on a variety of cell types. It is a potent immunosuppressant of macrophage functions. *In vitro*, IL-10 can inhibit the accessory function and antigen-presenting capacity of monocytes by, among other effects, downregulating class II MHC expression. Thus, IL-10 can inhibit monocyte/macrophage-dependent, antigen stimulated cytokine synthesis (especially IFN-γ) by human PBMNC and NK, and mouse Th1 cells. Additionally, IL-10 is a potent inhibitor of monocyte/macrophage activation and its resultant cytotoxic effects. As an immunostimulatory cytokine, IL-10 can act on B cells to enhance their viability, cell proliferation, Ig secretion, and class II MHC expression. Aside from B lymphocytes, IL-10 is also a growth co-stimulator for thymocytes and mast cells, as well as an enhancer of cytotoxic T cell development.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.