

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Dopamine β -Hydroxylase in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Dopamine β -Hydroxylase Leu37-Gly617 Accession # P09172
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

DBH (Dopamine β -Hydroxylase) is a 77-78 kDa member of the Cu⁺⁺ type II ascorbate-dependent monooxygenase family of enzymes. It is expressed in noradrenergic nerve terminals and adrenal medullary chromaffin cells, and serves as a catalyst for the conversion of dopamine into norepinephrine. Unlike its family member counterparts (TH and PNMT) that are cytosolic, DBH is embedded in the membranes of secretory vesicles. Human DBH is a 617 amino acid (aa) type II transmembrane glycoprotein (SwissProt # P09172). It contains an N-terminal 16 aa cytoplasmic region, plus a 580 aa luminal domain (aa 38-617). The luminal region possesses one DOMON (dopamine β -monooxygenase N-terminal) domain (aa 51-169) plus two consecutive monooxygenase motifs (aa 214-523). DBH exists as both a membrane-embedded 77-78 kDa isoform, and a soluble 73-75 kDa isoform. The latter may arise from cleavage following Gly39. Structurally, DBH will form a disulfide-linked homodimer, which then noncovalently associates with another DBH covalent homodimer, generating a functional 290 kDa homotetramer. Over aa 37-617, human DBH shares 79% aa sequence identity with mouse DBH.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.