

## DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse CD74 in direct ELISAs and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant human CD74 is observed.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant mouse CD74 Gln56-Leu215 Accession # NP_034675
<b>Conjugate</b>	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
<b>Formulation</b>	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

<b>Western Blot</b>	Optimal dilution of this antibody should be experimentally determined.
<b>Immunohistochemistry</b>	Optimal dilution of this antibody should be experimentally determined.

## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

## BACKGROUND

CD74 (also known as H2 class II histocompatibility antigen gamma chain and Ia antigen-associated invariant chain/Ii) is 31-41 kDa glycoprotein, member of no known protein family. It is expressed by multiple cell types, including dendritic cells, macrophages, immature and mature B cells, type II greater alveolar epithelial cells and colonic epithelium. CD74 is best known as an ER-embedded trimeric chaperone that directs MHC class II dimers through the endosome and into the cell membrane. When embedded in the endosome, it undergoes cleavage that both facilitates MHC dimer transit, and generates a cytosolic signaling fragment. It also appears on the cell surface where, in conjunction with CD44 (and possibly c-met), it binds MIF, inducing either cell proliferation, cell survival, or cytokine release. Mouse CD74 is a 279 amino acid (aa) type II transmembrane protein (SwissProt P04441). It possesses a 30 aa cytoplasmic region plus a 224 aa luminal/extracellular domain (aa 56-279) that contains one thyroglobulin type-1 repeat (aa 193-254). CD74 is known to be phosphorylated on Ser9, and occasionally undergo glycanation with chondroitin at Ser265 (on monocytes and B cells). There are multiple potential splice forms. The isoform used for antibody formation by R&D Systems, Inc. exhibits a deletion of aa 192-255 (GenBank NP\_034675). Two other splice forms show either the same deletion just described coupled to a four aa substitution for aa 277-279, or a 10 aa substitution for aa 131-279. Over aa 56-215 of the NP\_034675 isoform, mouse CD74 shares 86% and 69% aa sequence identity with rat and human CD74, respectively.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc. and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.