

DESCRIPTION


Species Reactivity	Rat
Specificity	Detects rat CD19 in direct ELISAs and Western blots. In direct ELISAs, approximately 25% cross-reactivity with recombinant human CD19 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line recombinant rat CD19 Arg19-Gly287 Accession # NP_001013255
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CD19 (also surface antigen B4 and Leu12) is a 95-110 kDa member of the Immunoglobulin superfamily of molecules. It is expressed by B cells, and interacts with CD21 for the purpose of reducing the threshold of the antigen signal needed to with activate the BCR. CD19 ligation also promotes B cell:follicular dendritic cell (FDC) interaction and B cell proliferation in the FDC zone of the spleen. Mature rat CD19 is a 529 amino acid (aa) type I transmembrane glycoprotein (aa 19-547). Based on mouse, it contains a 269 aa extracellular region (aa 19-287) plus a 236 aa cytoplasmic domain. The extracellular region possesses two C2-type Ig-like domains (aa 20-113 and 171-271) and one utilized phosphorylation site at Ser225. The cytoplasmic domain contains five potential Tyr phosphorylation sites. There is one splice form that shows a two aa substitution after Gly489. Over aa 19-287, rat CD19 shares 88% and 57% aa sequence identity with mouse and human CD19, respectively.