

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Desmocollin-2 in direct ELISAs. In direct ELISAs, approximately 10% cross-reactivity with recombinant human Desmocollin-2 is observed, and less than 1% cross-reactivity with recombinant mouse (rm) Desmocollin-1 and rmDesmocollin-3 is observed
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Desmocollin-2 Arg136-Pro694 Accession # P55292
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

DSC-2 (Desmocollin [Greek for "glue-that-binds"]-2; also epithelial type 2 desmocollin) is a 100-120 kDa member of the Ca⁺⁺-dependent cadherin family of adhesion molecules. It is found on the surface of both simple and stratified epithelium that form desmosomes. DSC-2 expression is induced by DSG-2 where it serves as a component of desmosomes, forming a linkage that unites adjacent cells with cytoplasmic intermediate filaments. Homodimeric DSC-2 forms in-trans, and DSC-2 may also form heterotypic interactions with DSG-2. A DSC-2:DSG-2 cis-interaction is suggested to block DSG-2 activation of Akt, an action that results in β-catenin mediated cell proliferation. Mature mouse DSC-2 is a 767 amino acid (aa) type I transmembrane glycoprotein (aa 136-902). The mature molecule contains a 559 aa extracellular region (aa 136-694) with five cadherin domains (aa 136-694) and a 187 aa cytoplasmic tail (aa 716-902). There is one 100-110 kDa splice variant that shows an 11 aa substitution for aa 838-902. Over aa 136-694, mouse DSC-2 shares 87% and 79% aa sequence identity with rat and human DSC-2, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.