

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human PAK1 in direct ELISAs. In direct ELISAs, less than 10% cross-reactivity with recombinant human (rh) PAK1B, rhPAK2, rhPAK4, rhPAK6, and rhPAK7 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human PAK1 Leu128-Gln242 Accession # Q13153
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

PAK1 (p21-activated kinase 1; also α -PAK and p65-PAK) is both a cytoplasmic and nuclear 58-60 kDa member of PAK group I, STE20 subfamily, STE Ser/Thr protein kinase family of molecules. It is widely expressed, and is upregulated in numerous cancers. PAK1 has multiple activities. It is responsible for regulating focal adhesions during motility. At the front edge of cells, PAK1 promotes focal adhesion formation by phosphorylating PIX, while at the back edge, it directs focal adhesion disassembly. PAK1 also impacts microtubule formation. Microtubules are composed of heterodimeric subunits containing an α - and β -tubulin polypeptide. The availability of α -tubulin for microtubule formation is based on the ability of PAK1 to phosphorylate a α -tubulin chaperone, tubulin cofactor B. PAK1 is activated following binding to active GTPases which induce PAK autophosphorylation. When active, PAK1 is a monomer; when inactive, it reportedly homodimerizes. Human PAK1 is 545 amino acids (aa) in length. It contains one GTPase binding domain (aa 75-105) plus a protein kinase catalytic region (aa 250-521). There are five SH3-binding motifs. Acetylation occurs on Lys256, and there are at least 14 potential phosphorylation sites, seven of which are known to be utilized. There is one splice variant that shows a 35 aa substitution for aa 518-545. Over aa 128-242, human PAK1 shares 96% aa sequence identity with mouse PAK1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.