

#### DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse Lymphotoxin- $\alpha$ /TNF- $\beta$ in direct ELISAs and Western blots. In direct ELISAs, 100% cross-reactivity with recombinant human Lymphotoxin- $\alpha$ /TNF- $\beta$ is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant mouse Lymphotoxin- $\alpha$ /TNF- $\beta$
<b>Conjugate</b>	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
<b>Formulation</b>	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

**Western Blot** Optimal dilution of this antibody should be experimentally determined.

#### PREPARATION AND STORAGE

**Shipping** The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

**Stability & Storage** Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

#### BACKGROUND

Tumor necrosis factor-beta (TNF- $\beta$ ), also known as lymphotoxin-alpha (LT- $\alpha$ ), is a secreted homotrimeric glycoprotein belonging to the TNF superfamily and is designated TNFSF1B. It is produced by NK, T, and B cells. TNF- $\beta$  was originally identified as protein that kills tumor cells in cell culture supernatants of a lymphoblastoid cell line. The TNF- $\beta$  subunit also associates with the type II transmembrane TNF superfamily protein lymphotoxin beta (LT $\beta$ ) to generate two types of heterotrimers designated as LT $\alpha$ 1 $\beta$ 2 (a single TNF- $\beta$  chain non-covalently associated with two chains of LT $\beta$ ), and LT $\alpha$ 2 $\beta$ 1 (1, 2). TNF- $\alpha$ , TNF- $\beta$ , and LT $\beta$  form a subfamily of the TNF related ligands. Their genes are genetically linked within a compact cluster inside the major histocompatibility complex locus (2, 3). The soluble TNF- $\beta$  binds and signals through TNF R1 and TNF R2. In contrast, the membrane-bound LT $\alpha$ 1 $\beta$ 2 interacts specifically with the LT $\beta$  receptor (LT $\beta$ R), which does not bind TNF- $\beta$  or TNF- $\alpha$ . Both TNFR1 and TNFR2 bind LT $\alpha$ 2 $\beta$ 1, which is recognized weakly by LT $\beta$  R (4, 5). TNF R1 and 2 express very broadly, while expression of LT $\beta$  R is restricted to stromal cells of lymphoid tissues. Herpesvirus entry mediator binds TNF- $\beta$  in vitro (6). The physiological importance of such interaction, if it occurs *in vivo*, is unclear. Distinct functions attributed to TNF- $\beta$  from transgenic knock-out mice include, loss of lymph node development, change in splenic architecture, impaired germinal center formation, and susceptibility to pulmonary tuberculosis (7, 8). TNF- $\beta$  also has overlapping physiological functions with LT $\beta$  and TNF- $\alpha$  in lymphoid organogenesis (7). Mouse and human TNF- $\beta$  share approximately 74% homology in their amino acid sequence.

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.