

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Cathepsin H in direct ELISAs and Western blots. In direct ELISAs, approximately 15% cross-reactivity with recombinant mouse Cathepsin H is observed, and less than 1% cross-reactivity with recombinant human Cathepsin L2 is observed.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human Cathepsin H Ala23-Val335 Accession # CAA34734
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.2 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.2 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Cathepsin H (CTSH; alos CPSB) is a 28-30 kDa ubiquitously expressed member of the peptidase C1 family of cysteine lysosomal enzymes. Depending upon the isoform, it may either be secreted or retained in the lysosomes. Although this family of enzymes plays a key role in general intracellular molecule degradation, CTSH is also known to participate in other activities. Such activities include the activation of progranzyme B, the late stage processing of pulmonary surfactant protein B, and the cleavage, in endosomes, of TLR3, allowing for subsequent TLR signaling. Human proCTSH is 335 amino acids (aa) in length. It contains one 22 aa signal peptide, a 75 aa prosequence (aa 23-97), an intervening "mini fragment" (aa 98-105), a second prosequence (aa 106-115), and a 220 aa C-terminal mature region that could be further cleaved after Asn292 into two peptide fragments. CTSH is self-activating. The 40-42 kDa proprecursor (aa 23-335) is cleaved thrice to initially generate a 30 kDa intermediate form, followed by a 28 kDa mature form that is covalently attached to the aforementioned "mini fragment". Further processing at the C-terminus may create a smaller 24-25 kDa mature segment. CTSH is unusual in that it possess both aminopeptidase and endopeptidase activity. While endopeptidase activity is intrinsic to the mature region, aminopeptidase activity requires the presence of a covalently-linked "mini fragment". There is one splice variant that shows a deletion of aa 11-22. Although this occurs within the signal sequence, the molecule is secreted and shows high enzymatic activity. Over aa 23-335, human CTSH shares 84% aa sequence identity with mouse CTSH.