

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse, and rat CIB1 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CIB1 Gly2-Leu191 Accession # Q99828
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CIB1 (Calcium and Integrin-Binding protein 1; also calmyrin, KIP and CIBP) is a 22-23 kDa, Ca⁺⁺-binding member of the CIB family of proteins. It is widely expressed, being found in neurons, platelets/megakaryocytes, skeletal muscle myocytes and lymphocytes. CIB1 is associated with the cytosolic side of the plasma membrane, and has multiple binding partners, including InsP₃R, GPIIb/αIIb, presenilin 2, and NBR1 plus FEZ. When CIB1 binds InsP₃R, this ER-embedded receptor is both activated, and later desensitized to subsequent ligand binding. Relative to GPIIb, CIB1 interaction with the αIIbβ₃ integrin on platelets following thrombin exposure appears to inhibit integrin activation, thus providing a tight control on subsequent platelet binding to fibrinogen. Human CIB1 is 191 amino acids (aa) in length. It contains a utilized myristoylation site at Gly2, followed by two EF-hand domains (aa 103-183). CIB1 is suggested to act as a monomer. There is at least one potential isoform variant that shows a 40 aa insertion after Lys29. Full-length human CIB1 (aa 1-191) shares 94% aa sequence identity with mouse CIB1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.