

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse NAIP in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant mouse NAIP2 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse NAIP Lys783-Ser856 Accession # Q9QWK5
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

NAIP-1 (Neuronal Apoptosis Inhibitory Protein 1; also Naip and Birc1) is a 150-160 kDa member of both the NLR/PRR and IAP family of proteins. It is widely, but not ubiquitously expressed, and is found in adipocytes, macrophages, neurons, renal proximal and distal convoluted tubules, and nonproliferative intestinal epithelium. NAIP-1 plays a key regulatory role in caspase activation. It promotes caspase-1 activation, blocks caspase-3 and -7 activation, and by binding neuronal hippocalcin, inhibits caspase-3-independent apoptosis. Mouse NAIP-1 is 1403 amino acids (aa) in length. It contains three consecutive N-terminal BIR (Baculovirus Inhibition of apoptosis Repeat) domains (aa 63-346) and one NACHT region (aa 464-758). The BIR #2 domain is posited to bind both caspase-3 and -7, while the BIR domain #3 likely interacts with hippocalcin. NAIP-1 undergoes proteolysis, generating an N-terminal 100 kDa fragment (likely aa 1- ~ 869) and a 60 kDa C-terminal fragment (likely aa 870-1403). The C-terminal fragment undergoes further processing to generate a 25 kDa and 35 kDa fragment. There are at least seven numbered NAIP genes in mouse that have arisen by gene duplication. Over the aa range 783-856, they share less than 60% aa sequence identity with NAIP-1. Over the same aa range, mouse NAIP-1 shares 54% aa sequence identity with the only human NAIP gene.