

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Glis1 in direct ELISAs and Western blots. In direct ELISAs, approximately 60% cross-reactivity with recombinant mouse Glis1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Glis1 Met446-Thr620 Accession # Q5VTL4
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Glis1 (GLI/glioma-Similar 1; also GliH1) is a 66 kDa (predicted) member of the Kruppel-like zinc finger family of proteins. In the adult, it is principally expressed in placenta and kidney, while in the embryo, it has widespread expression in mesodermal structures. Glis1 can be induced in suprabasal keratinocytes by activating cytokines, and is believed to promote a unique differentiation program. Glis1 is also effective in generating iPSCs from somatic cells (fibroblasts) when concurrently expressed with Oct3, Sox2 and Klf4. Multiple reprogramming pathways are involved, and include Myc, Wnt and Nanog molecules. Functionally, Glis1 is believed to act as a transcriptional repressor. Although it has been described as possessing an activation domain, it would appear that gene promotion is a consequence of the repression of a "repressor". The consequence sequence ACCACCCA corresponds to a GLIS binding motif. Human Glis1 is 620 amino acids (aa) in length. Based on mouse, it contains a zinc-finger repressor region (aa 197-346) plus a putative transactivation domain (aa 448-620). And again based on mouse, there might be an approximately 796 aa splice form that shows an alternative start site 171 aa upstream of the 620 aa isoform. The existence of such an isoform would generate an approximately 95-100 kDa isoform in SDS-PAGE. Over aa 446-620, human Glis1 shares 80% aa sequence identity with mouse Glis1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.