

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TIGAR/C12orf5 in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human TIGAR/C12orf5 Ala2-Arg270 Accession # Q9NQ88
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide

*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

TIGAR (TP53-Inducible Glycolysis and Apoptosis Regulator; also C12orf5) is a 29-30 kDa member of the phosphoglycerate mutase family of molecules. TIGAR is likely widely expressed, and serves as a target of p53 activity. Cellular glucose is normally converted into Glu-6-P. This compound can either be converted into Fru-6-P for use in ATP production, or directed into the pentose phosphate to create NADPH. If Fru-6-P is formed, it can be acted upon by either PFK1 or PFK2. PFK1 creates Fru-1,6-P2, which is subsequently broken down into pyruvate for use in the Krebs cycle. PFK2 creates Fru-2,6-P2, which is not an energy source but a regulator that promotes the activity of PFK1. TIGAR dephosphorylates Fru-2,6-P2, creating Fru-6-P. This both inhibits PFK1 activity, and drives the system backwards, recreating Glu-6-P which enters the pentose phosphate shunt and generates NADPH. p53 induces TIGAR synthesis, with a resulting increase in NADPH. NADPH increases glutathione which neutralizes ROS, and blocks activation of caspase-2. The emphasis away from energy production also leads to the activation of other pathways that produce molecules involved in DNA repair. Human TIGAR is 270 amino acids (aa) in length. It contains one histidine phosphatase domain (aa 6-90) plus two utilized phosphorylation sites. There is one potential alternative start site at Met60. Full-length human TIGAR shares 72% aa sequence identity with mouse TIGAR.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.