

#### DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human ALK-7 in direct ELISAs.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human ALK-7 Met1-Glu113 Accession # Q8NER5
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Immunohistochemistry</b>	10-25 µg/mL	See Below

#### DATA

##### Immunohistochemistry



**ALK-7 in Human Breast Cancer Tissue.**  
ALK-7 was detected in immersion fixed paraffin-embedded sections of human breast cancer tissue using Sheep Anti-Human ALK-7 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF7749) at 15 µg/mL for 1 hour at room temperature followed by incubation with the Anti-Sheep IgG VisUCyte™ HRP Polymer Antibody (Catalog # VC006). Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using DAB (brown) and counterstained with hematoxylin (blue). Specific staining was localized to cell surfaces in cancer cells. View our protocol for [IHC Staining with VisUCyte HRP Polymer Detection Reagents](#).

#### PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

#### BACKGROUND

Activin receptor-like kinase 7 (ALK-7), also known as Activin R1C (gene name ACVR1C), is a glycosylated 58 kDa type I receptor in the superfamily of TGF-β serine/threonine kinase receptors. It associates with type II receptors to form a signaling complex that responds to the ligands Activin AB, and Activin B, GDF3, and Nodal. ALK-7 plays a role in regulating energy balance by inhibiting insulin secretion and inducing pancreatic beta cell apoptosis. It is expressed in adipose tissue but downregulated in obesity. ALK-7 is also expressed in pituitary gonadotropic cells and in pre-eclamptic placenta. It induces the apoptosis of trophoblasts as well as ovarian granulosa and epithelial cells. Within the extracellular domain, human ALK-7 shares 95% and 91% amino acid (aa) sequence identity with mouse and rat ALK-7, respectively. Alternate splicing of human ALK-7 generates additional isoforms with either a 50 aa N-terminal truncation or with deletions of 79 aa or 157 aa that encompass the transmembrane segment.