

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse SECTM1A in direct ELISAs.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse SECTM1A Met1-Thr165 Accession # Q921W8
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Blockade of Receptor-ligand Interaction	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SECTM1A (Secreted and Transmembrane 1A), is 192 amino acid (aa) protein that shares approximately 45-51%, 81%, and 43% aa sequence identity with human SECTM1 (also called K12), rat SECTM1, and mouse SECTM1B/K12, respectively, and appears to share structural and functional characteristics with other SECTM1 proteins. Human SECTM1 can be found either found as an approximately 27 kDa intracellular type I transmembrane protein that shows a perinuclear, Golgi-like staining pattern, or as a 20 kDa soluble, secreted form, and is produced by some myeloid cells and by thymic epithelia and fibroblasts (1-3). Stimulation with IFN-γ is often necessary to detect human SECTM1 expression, and it is thought to be an interferon early-response gene (1-5). Mouse SECTM1A cDNA encodes a signal sequence, an extracellular domain with four potential N-linked glycosylation sites, a transmembrane sequence, and a very short (approximately 6 aa) cytoplasmic sequence (6). SECTM1 proteins from human and mouse show species-specific binding of CD7 and co-stimulation of T cells, including enhancement of CD3-induced proliferation (2-4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.