

DESCRIPTION

| | |
|---------------------------|---|
| Species Reactivity | Mouse |
| Specificity | Detects mouse Thrombospondin-4 in direct ELISAs and Western blots. In direct ELISAs, approximately 10% cross-reactivity with recombinant human Thrombospondin-4 is observed, and less than 1% cross-reactivity with recombinant mouse Thrombospondin-1 is observed. |
| Source | Polyclonal Sheep IgG |
| Purification | Antigen Affinity-purified |
| Immunogen | Chinese hamster ovary cell line CHO-derived recombinant mouse Thrombospondin-4 Gln27-Asn963 (predicted) Accession # Q9Z1T2 |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS. |


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------------|---------------------------|-----------|
| Western Blot | 1 µg/mL | See Below |
| Immunohistochemistry | 5-15 µg/mL | See Below |


DATA

Western Blot



Detection of Mouse Thrombospondin-4 by Western Blot. Western blot shows lysates of mouse muscle tissue. PVDF membrane was probed with 1 µg/mL of Sheep Anti-Mouse Thrombospondin-4 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF7860) followed by HRP-conjugated Anti-Sheep IgG Secondary Antibody (Catalog # HAF016). A specific band was detected for Thrombospondin-4 at approximately 120 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.

Immunohistochemistry



Thrombospondin-4 in Mouse Skeletal Muscle. Thrombospondin-4 was detected in perfusion fixed frozen sections of mouse skeletal muscle using Sheep Anti-Mouse Thrombospondin-4 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF7860) at 5 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Sheep HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS019) and counterstained with hematoxylin (blue). Specific staining was localized to connective tissue. View our protocol for [Chromogenic IHC Staining of Frozen Tissue Sections](#).

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Reconstitution | Reconstitute at 0.2 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

BACKGROUND

Mouse Thrombospondin-4 (TSP-4) is a 120-130 kDa adhesive glycoprotein that belongs to the thrombospondin family of proteins. It is 963 amino acids (aa) in length and highly modular in structure, containing one Lamnin G-like domain, four EGF-like domains and eight type 3 TSP domains. It circulates as a disulfide-linked homopentamer, and is up-regulated in the heart in response to ischemic injury and pathology. Mature mouse TSP-4 is 93% aa identical to human TSP4.