

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human NQO-2 in direct ELISAs and Western blots. In direct ELISAs, less than 3% cross-reactivity with recombinant human NQO-1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human NQO-2 Ala2-Gln231 (Leu47Phe) Accession # P16083
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

NQO-2 (NADPH Quinone acceptor Oxidoreductase 2; also QR2, melatonin receptor 3/MT3 and NRH dehydrogenase) is a 25-26 kDa cytoplasmic flavoprotein member of the NAD(P)H dehydrogenase family of enzymes. It shows restricted expression, being found in retinal pigment epithelium, prostatic fibroblasts, select CNS neurons and RBCs. Like NQO-1, NQO-2 would appear to reduce catechol quinones, but without the use of NADH. In contrast to NQO-1, NQO-2 would appear to activate some quinones, generating cytotoxic products. NQO-2 also reportedly binds and stabilizes key cell protection molecules such as p53, thus acting as a gatekeeper for proteasome-mediated protein turnover. Finally, NQO-2 reportedly binds to melatonin, an act that may promote the subsequent binding of the androgen receptor and an inhibition of cell proliferation. Human NQO-2 is 231 amino acids (aa) in length. It contains a flavodoxin-like domain (aa 4-212) and three utilized Ser phosphorylation sites. NQO-2 functions as a noncovalent homodimer. There are two potential isoform variants. One shows an alternative start site at Met117, while another shows a deletion of aa 102-139. Full length human NQO-2 shares 81% aa sequence identity with mouse NQO-2, and less than 50% aa sequence identity with human NQO-1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.