

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse Doc2α in Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human Doc2β is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Doc2α Met1-Lys114 (Gly48Ser) Accession # Q14183
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

DOC2A/Doc2α (Double C2-like domain containing protein alpha) is a 44 kDa (predicted) monomeric member of the C2 domain-containing protein family of molecules. It is expressed in both neurons and mast cells, and appears to serve as an intracellular Ca⁺⁺ sensor protein that regulates secretory vesicle release. In neurons, Doc2α is normally bound to synaptic vesicles and interacts with Munc13-1 to promote secretory vesicle exocytosis through the cell membrane. In mast cells, a similar process occurs that involves Munc13-4 instead of Munc13-1. Human Doc2α is 400 amino acids (aa) in length. It contains a Mid domain (aa 13-37) that binds Munc13-1, followed by one C2 domain that binds Ca⁺⁺ and lipid (aa 91-195), and a second C2 domain that binds SNAP25 (253-356). There is one potential alternative start site 16 aa upstream of the standard site. Over aa 1-114, human Doc2α shares 90% aa sequence identity with mouse Doc2α. Human DOC2B is the product of a separate gene, and shares no meaningful aa sequence identity (< 30%) with human Doc2α.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.