

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Dihydrofolate Reductase/DHFR in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Dihydrofolate Reductase/DHFR Met1-Asp187 Accession # P00374
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

DHFR (DiHydroFolate Reductase; also Tetrahydrofolate dehydrogenase) is a 21-23 kDa member of the dihydrofolate reductase family of enzymes. It is a ubiquitously expressed monomer, and considered to be a housekeeping gene. Housekeeping genes are those that play a role in multiple pathways, although not the same pathway(s) in all cells. DHFR participates in the reduction of dihydrofolate to tetrahydrofolate, a product that is subsequently used in the synthesis of purines and thymidylic acid that are used to generate both RNA and DNA. Within the cell, DHFR is known to exist in two pools: one contains DHFR bound to its own RNA where it acts as a transcriptional repressor, while another contains DHFR bound to NADPH. Human DHFR is 187 amino acids (aa) in length and possesses one DHFR domain (aa 4-185). Its mRNA binding motif is suggested to involve Cys6, Leu22, Glu30 and Ser118. There is one potential alternative start site found 75 aa upstream of the standard start site. Full length human DHFR (aa 1-187) shares 90% aa sequence identity with mouse DHFR.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.