

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Piccolo in direct ELISAs.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Piccolo Ala4065-Phe4328 Accession # NP_149015
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Piccolo (PCLO; also known as Aczonin) is a 530-570 kDa member of a small family of zinc finger proteins that are associated with the cytoskeletal matrix of exocytosis-competent cells. Its name (as well as that of oboe and bassoon) derives from the assumption that this molecule is part of an ensemble of "presynaptic" proteins that orchestrate secretory events at (presynaptic) cell membranes. Piccolo is known to be initially present in the trans-Golgi of neuronal cells. When needed, it is transported to the presynaptic density (or active zones) of GABAergic and glutamergic neurons, and to the plasma membrane of pancreatic β -cells. It reportedly forms homodimers and heterodimers with Rim2, and serves as a Ca^{2+} sensor for exocytosis. Human Piccolo is 5065 amino acids (aa) in length. It is a multidomain protein with N-terminal Pro- and Gln-rich sequences, two C4-type Zn-finger motifs (aa 535-1028), one PZD domain (aa 4427-4478) and two membrane-binding C2 domains (aa 4565-5057). There are at least 30 potential Thr/Ser and two Tyr phosphorylation sites. Multiple splice variants exist that principally involve insertions and deletions. None of the splice forms are missing, or show substitutions for, the range of amino acids used by R&D Systems for antibody production. Two splice forms contain identical insertions of 7, 8, 9, 17, 43 and 54 aa, and differ only in a deletion of 60 aa at the C-terminus. A third splice variant shows a 48 aa substitution for aa 443-5065, while a fourth isoform utilizes an alternative start site at Met4459 that is coupled to 1) a 17 aa insertion after Lys4570, and 2) a deletion of aa 4798-5065. Over aa 4065-4328, human Piccolo shares 97% aa sequence identity with mouse Piccolo.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.