

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat LYVE-1 in direct ELISA and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant mouse LYVE-1 is observed, and approximately 10% cross-reactivity with recombinant human LYVE-1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant rat LYVE-1 Thr53-Thr259 Accession # NP_001099756
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

LYVE-1 (Lymphatic vessel endothelial hyaluronic acid receptor 1; also CRSBP-1) is a 58-64 kDa, monomeric, glycoprotein member of the Link protein superfamily of hyaluron-binding molecules. It has limited expression, being found on the cell surface of lymphatic endothelial cells, endothelial cells of lymphoid sinuses, nodal stromal cells, and macrophages plus dendritic cells. HA (hyaluronan) is a nonsulfated, freestanding, repeating disaccharide consisting of GlcA (glucuronic acid) in β -linkage with GlcNAc (N-acetylglucosamine). It should not be confused with heparan, which is sulfated, protein-linked, and composed of repeating GlcA/IdoA and GlcNAc residues in both α - and β -linkages. HA is ubiquitous and occupies space between collagen fibers. It undergoes both normal, and pathology-induced turnover, and presumably does so by binding to LYVE-1, CD44 and HARE on lymphatic endothelium. Ultimately, HA is transported to the liver and nodes where it undergoes degradation. This may be necessary as low MW HA is proinflammatory. LYVE-1 is also a receptor for PDGF-BB and VEGF-A, and LYVE-1 ligation apparently induces endothelial cell contraction with the opening of intercellular junctions. Based on mouse, rat LYVE-1 is synthesized as a 343 amino acid (aa) type I transmembrane protein that contains a very long signal sequence (aa 1-52). The extracellular region is likely to be 182 aa in length (aa 53-234) and contain one Link domain (aa 62-210). LYVE-12 is likely maintained in a default "off mode" by undergoing sialylation, possibly at Thr83. Over aa 53-259, rat LYVE-1 shares 82% and 60% aa sequence identity with mouse and human LYVE-1, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.