

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CXCL4/PF4 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CXCL4/PF4 Glu32-Ser101 Accession # P02776
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CXCL4, also known as Platelet Factor 4, is a member of the CXC chemokine family, CXCL4. Human CXCL4 is a 101 amino acid (aa) protein with a 32 aa signal sequence and a 70 aa mature protein. CXCL4 has homology with IL-8 and β -thromboglobulin. Human and mouse CXCL4 share a 64% identity. CXCL4 contains several heparin-binding sites at the C-terminal region. The active protein consists of a tetramer composed of individual CXCL4 subunits. Megakaryocytes synthesize CXCL4 and store it as tetramers in α -granules. The CXCL4 tetramers are secreted by activated platelets and can be measured at micromolar levels in serum. In contrast to other CXC chemokines, CXCL4 lacks chemotactic activity for polymorphonuclear granulocytes. CXCL4 does not contain an ELR motif. However, many other functions have been observed for CXCL4. CXCL4 is involved in monocyte survival and differentiation into macrophages, and it has anti-angiogenic activity. CXCL4 has been demonstrated to inhibit the binding of FGF-2 to high-affinity receptors and its subsequent internalization. Cell surface neutrophil chondroitin sulfate chains serve as CXCL4 binding sites; affinity is controlled by the degree of sulfation of these chains.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.