

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse and rat in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human NADK Met1-Gly446 Accession # O95544
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

NADK (Nicotinamide Adenine Dinucleotide Kinase; also Poly(P)/ATP NAD kinase) is a 49-52 kDa cytosolic member of the NAD kinase family of enzymes. Although it is widely expressed, it is not found in skeletal muscle. NADK forms homotetramers and catalyzes the generation of NADP from NAD through the hydrolysis of ATP. NADP plays a role in fatty acid synthesis plus steroidogenesis, and provides a substrate for the generation of NADPH that serves to regulate the expression of reductases and antioxidant proteins. Human NADK is 446 amino acids (aa) in length. It contains one kinase domain (aa 105-429) plus a C-terminal Ala- and Glu-rich segment. There are four utilized phosphorylation sites at Ser46, Ser48, Ser50 and Ser64. The potential exists for an alternate start site at Met134, and two isoform variants are reported, one that contain a 56 aa substitution for aa 1-88, and another that possesses a 105 aa substitution for Met88 coupled to a 41 aa insertion after Glu131. Full-length NADK (aa 1-446) shares 91% aa sequence identity with mouse NADK. Recently, a human mitochondrial NADK was identified (termed C5orf33), but found to share only 19% aa sequence identity with cytosolic NADK.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.