

DESCRIPTION




Species Reactivity	Mouse/Rat
Specificity	Detects mouse and rat in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse Integrin $\beta 5$ Gly24-Asn719 Accession # NP_001139356
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 μ g/mL	See Below
Flow Cytometry	2.5 μ g/10 ⁶ cells	See Below
Immunohistochemistry	5-15 μ g/mL	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

The β 5 integrin is a 95-110 kDa member of the integrin beta chain family, integrin superfamily of molecules. It is expressed by select cell types, including microglia, keratinocytes, fibroblasts, endothelial cells, adhesive monocytes and macrophages, retinal pigment epithelium, immature dendritic cells and colonic epithelium. The Integrin β 5 subunit forms a noncovalent heterodimer with the α integrin subunit, and together they bind vitronectin, and the Thy-1 antigen. α β 5 is reported to participate in phagocytosis, and act in concert with CD81, CD36 and stabilin-2 on the cell surface. When α β 5 binds Thy-1, the heterodimer apparently interferes with cell membrane activation of TGF- β . Intracellularly, it appears to regulate actin skeleton remodeling via focal adhesion complex formation, and may contribute to cryptic pathways associated with antigen presentation. Mature mouse Integrin β 5 is a 775 amino acid (aa) type I transmembrane glycoprotein that contains a 696 aa extracellular domain (ECD) (aa 24-719) and a 56 aa cytoplasmic region. The ECD contains a 243 aa vWF-A domain (aa 136-378) and a Cys-rich tandem repeat region (aa 465-630). There is one splice variant that shows a 57 aa substitution for aa 760-798. This splice form does not appear to be expressed by the human Integrin β 5 gene. Over aa 24-719, mouse Integrin β 5 shares 91% and 97% aa sequence identity with human and rat Integrin β 5, respectively.